



A mixed problem for a Boussinesq hyperbolic equation with integral condition

Said Mesloub, Abdelouahab Mansour

► To cite this version:

Said Mesloub, Abdelouahab Mansour. A mixed problem for a Boussinesq hyperbolic equation with integral condition. 2005. hal-00338641

HAL Id: hal-00338641

<https://hal.science/hal-00338641>

Submitted on 14 Nov 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A MIXED PROBLEM FOR A BOUSSINESQ HYPERBOLIC EQUATION WITH INTEGRAL CONDITION

SAID MESLOUB (1) AND A. MANSOUR (2)

ABSTRACT. A hyperbolic problem wich combines a classical(Dirichlet) and a non-local contraint is considered.The existence and uniqueness of strong solutions are proved,we use a fonctionnal analysis method based on a priori estimate and on the density of the range of the operator generated by the considered problem.

1.INTRODUCTION

The first study of evolution problems with a nonlocal condition - the so called energy specification - goes back to Cannon[5] ,1963 Using an integral condition ,we proved the existence and uniqueness of the solution of a mixed problem wich combine a classical (Dirichlet)and an integral condition for the equation . Problems involving local and integral condition for hyperbolic equations are investigated by the energy inequalities method in [1] , [6] , [7] , [8] , [9] , [10] , [11], [12] .In this paper ,we prove the existence and uniqueness of the solution for the mixed problem (1) – (5) .Our proof is based on a priori estimate and on the fact that the range of the operator generated by the considered problem is dense.

Date: Octobre 30, 2008.

Key words and phrases. hyperbolic equation, integral condition, a priori estimate.

2. Formulation of the problem

In the region $Q = (0, l) \times (0, T)$, with $l < \infty$ and $T < \infty$, we shall consider the problem

$$(1) \quad Lu = u_{tt} - (b(x, t)u_x)_x - \beta \frac{\partial^4 u}{\partial t^2 \partial x^2} = f(x, t), \forall (x, t) \in Q$$

$$(2) \quad l_1 u = u(x, 0) = \varphi_1(x), \quad x \in (0, l)$$

$$(3) \quad l_2 u = u_t(x, 0) = \varphi_2(x), \quad x \in (0, l)$$

$$(4) \quad u(0, t) = 0, \quad t \in (0, T)$$

$$(5) \quad \int_0^l x u(x, t) dx = 0, \quad t \in (0, T)$$

where $\beta \in IR_+^*$ and $b(x, t)$ and its derivatives satisfy the conditions:

$$\begin{aligned} C_1 : & b_0 \leq b(x, t) \leq b_1, \quad b_t(x, t) \leq b_2, \quad b_x(x, t) \leq b_3, \\ & \text{for any } (x, t) \in \overline{Q}, \\ C_2 : & b_{tt}(x, t) \leq b_4, \quad b_{xt}(x, t) \leq b_5, \text{ for any } (x, t) \in \overline{Q}. \end{aligned}$$

The functions f , φ_1 and φ_2 are known functions which satisfy the compatibility conditions:

$$\varphi_1(0) = \varphi_2(0) = \int_0^l x \varphi_1(x) dx = \int_0^l x \varphi_2(x) dx = 0.$$

3. functional Spaces

The problem (1)-(5) can be put in the following operator form: $Lu = \mathcal{F}$, $u \in D(L)$, where:

$$Lu = (\mathcal{L}u, l_1u, l_2u) \text{ and } \mathcal{F} = (f, \varphi_1, \varphi_2).$$

The operator L is considered from B to H , where B is the Banach space consisting of functions $u \in L^2(Q)$, satisfying conditions (4) and (5) with the finite norm:

$$\|u\|_B^2 = \sup_{0 \leq \tau \leq T} \left[\|u(\cdot, \tau)\|_{L^2(0,l)}^2 + \|u_t(\cdot, \tau)\|_{L^2(0,l)}^2 \right]$$

and F is the Hilbert space $L^2(Q) \times L^2(0, l) \times L^2(0, l)$ equipped with the norm:

$$\|\mathcal{F}\|_H^2 = \|f\|_{L^2(Q^\tau)}^2 + \|\varphi_1\|_{L^2(0,l)}^2 + \|\varphi_2\|_{L^2(0,l)}^2.$$

Let $D(L)$ denote the domain of L , which is the set of all functions $u \in L^2(Q)$ for which $u_t, u_x, u_{tx}, u_{tt}, u_{ttx} \in L^2(Q)$ and satisfying conditions (4) and (5).

4. A priori estimate and its consequences

Theorem 1: For any function $u \in D(L)$ satisfies conditions C₁-C₂ there exists a positive constant c , such that

$$(7) \quad \|u\|_B \leq c \|Lu\|_H,$$

Proof : We consider the scalar product in $L^2(Q^\tau)$ of the operator $\mathcal{L}u$ and Mu , where $Mu = x\mathfrak{S}_x^*u_t - \mathfrak{S}x^*(\rho u_t)$, with $Q^\tau = (0, l) \times (0, \tau)$, $0 \leq \tau \leq T$, and $\mathfrak{S}_x^*v = \int_x^l v(\xi, t)d\xi$, we obtain

$$\begin{aligned}
(8) \quad (\mathcal{L}u, Mu)_{L^2(Q^\tau)} &= (u_{tt}, x\mathfrak{S}_x^*u_t)_{L^2(Q^\tau)} - ((b(x, t)u_x)_x, x\mathfrak{S}_x^*u_t)_{L^2(Q^\tau)} \\
&\quad - \beta(u_{ttxx}, x\mathfrak{S}_x^*u_t)_{L^2(Q^\tau)} - (u_{tt}, \mathfrak{S}_x^*(\rho u_t))_{L^2(Q^\tau)} \\
&\quad + ((b(x, t)u_x)_x, \mathfrak{S}_x^*(\rho u_t))_{L^2(Q^\tau)} + \beta(u_{ttxx}, \mathfrak{S}_x^*(\rho u_t))_{L^2(Q^\tau)}.
\end{aligned}$$

Making use of conditions (2)-(5) and integrating by parts we establish the equalities:

$$\begin{aligned}
(9) \quad (u_{tt}, x\mathfrak{S}_x^*u_t)_{L^2(Q^\tau)} &= \frac{1}{2} \|\mathfrak{S}_x^*u_t(\cdot, \tau)\|_{L^2(0, l)}^2 \\
&\quad - \frac{1}{2} \|\mathfrak{S}_x^*\varphi_2\|_{L^2(0, l)}^2 - (\mathfrak{S}_x^*u_{tt}, u_t)_{L_\rho^2(Q^\tau)},
\end{aligned}$$

$$\begin{aligned}
(10) \quad &- ((b(x, t)u_x)_x, x\mathfrak{S}_x^*(u_t))_{L^2(Q^\tau)} \\
&= \frac{1}{2} \left\| \sqrt{b(\cdot, \tau)}u(\cdot, \tau) \right\|_{L^2(0, l)}^2 - \frac{1}{2} \left\| \sqrt{b(\cdot, 0)}\varphi_1 \right\|_{L^2(0, l)}^2 \\
&\quad - \frac{1}{2} \left\| \sqrt{b_t(\cdot, t)}u \right\|_{L^2(Q^\tau)}^2 - (b_x(x, t)u, \mathfrak{S}_x^*u_t)_{L^2(Q^\tau)} \\
&\quad - (b(x, t)u_x, u_t)_{L_\rho^2(Q^\tau)},
\end{aligned}$$

$$\begin{aligned}
(11) \quad -\beta(u_{ttxx}, x\mathfrak{S}_x^*(u_t))_{L^2(Q^\tau)} &= \frac{\beta}{2} \|u_t(\cdot, \tau)\|_{L^2(0, l)}^2 \\
&\quad - \frac{\beta}{2} \|\varphi_2\|_{L^2(0, l)}^2 - \beta(u_{ttx}, u_t)_{L_\rho^2(Q^\tau)}.
\end{aligned}$$

$$(12) \quad - (u_{tt}, \mathfrak{S}_x^*(\rho u_t))_{L^2(Q^\tau)} = (\mathfrak{S}_x^*u_{tt}, u_t)_{L_\rho^2(Q^\tau)},$$

$$(13) \quad ((b(x, t)u_x)_x, \mathfrak{S}_x^*(\rho u_t))_{L^2(Q^\tau)} = (b(x, t)u_x, u_t)_{L_\rho^2(Q^\tau)},$$

$$(14) \quad \beta(u_{ttxx}, \mathfrak{S}_x^*(\rho u_t))_{L^2(Q^\tau)} = \beta(u_{ttx}, u_t)_{L_\rho^2(Q^\tau)}.$$

Combining equalities (9)-(14) and (8) we obtain :

$$\begin{aligned}
 (15) \quad & \frac{1}{2} \|\mathfrak{S}_x^* u_t(\cdot, \tau)\|_{L^2(0,l)}^2 + \frac{1}{2} \left\| \sqrt{b(\cdot, t)} u(\cdot, \tau) \right\|_{L^2(0,l)}^2 \\
 & + \frac{\beta}{2} \|u_t(\cdot, \tau)\|_{L^2(0,l)}^2 \\
 & = \frac{1}{2} \|\mathfrak{S}_x^* \varphi_2\|_{L^2(0,l)}^2 + \frac{1}{2} \left\| \sqrt{b(\cdot, t)} \varphi_1 \right\|_{L^2(0,l)}^2 \\
 & + \frac{\beta}{2} \|\varphi_2\|_{L^2(0,l)}^2 + \frac{1}{2} \left\| \sqrt{b_t} u \right\|_{L^2(Q^\tau)}^2 + (b_x(x, t) u, \mathfrak{S}_x^* u_t)_{L^2(Q^\tau)} \\
 & + (\mathcal{L}u, x \mathfrak{S}_x^* u_t)_{L^2(Q^\tau)} - (\mathcal{L}u, \mathfrak{S}_x^*(\rho u_t))_{L^2(Q^\tau)}.
 \end{aligned}$$

By applying the Cauchy inequality to the last three terms on the right-hand side of the inequality (15) and making use conditions C₁, combining with (15), we obtain

$$\begin{aligned}
 (16) \quad & \|u(\cdot, \tau)\|_{L^2(0,l)}^2 + \|u_t(\cdot, \tau)\|_{L^2(0,l)}^2 \\
 & + \|\mathfrak{S}_x^* u_t(\cdot, \tau)\|_{L^2(0,l)}^2 \\
 & \leq k \left[\|f\|_{L^2(Q^\tau)}^2 + \|\varphi_1\|_{L^2(0,l)}^2 + \|\varphi_2\|_{L^2(0,l)}^2 \right. \\
 & \quad \left. \|u\|_{L^2(Q^\tau)}^2 + \|u_t\|_{L^2(Q^\tau)}^2 + \|\mathfrak{S}_x^* u_t\|_{L^2(Q^\tau)}^2 \right].
 \end{aligned}$$

where $k = \frac{\max(2, b_1, \beta + l^2, b_3^2 + b_2, l^4)}{\min(1, b_0, \beta)}$.

Applying the Gronwall lemma to (16), and eliminating the term $\|\mathfrak{S}_x^* u_t(\cdot, \tau)\|_{L^2(0,l)}^2$ of the left-hand side of the inequality we obtain

$$\begin{aligned}
 (17) \quad & \|u(\cdot, \tau)\|_{L^2(0,l)}^2 + \|u_t(\cdot, \tau)\|_{L^2(0,l)}^2 \\
 & \leq k \exp(kT) \left(\|f\|_{L^2(Q^\tau)}^2 + \|\varphi_1\|_{L^2(0,l)}^2 + \|\varphi_2\|_{L^2(0,l)}^2 \right).
 \end{aligned}$$

Since the left-hand side of (17) does not depend on τ , we take the supremum with τ from 0 to T , then the estimate (7) follows with $c = \sqrt{k} \exp(k\frac{T}{2})$.

5. Solvability of the problem

Proposition 1. The operator L acting from B to H have a closure.

Proof. (see [3])

Let be \bar{L} the closure of L , $D(\bar{L})$ its domain .

Definition . The solution of $\bar{L}u = F$ for any $u \in D(\bar{L})$ is strong solution of problem(1)-(5). we take the limit in the inequality (7) ,we obtain $\|u\|_B \leq c \|\bar{L}u\|_H, \forall u \in D(\bar{L})$.

From the inequality we have:

Corollary 1: The strong solution of problem (1)-(5) when it exists, it's unique, and depends continuly of data f, φ_1, φ_2 .

Corollary 2 : The set of values $R(\bar{L})$ of the operator \bar{L} is equal to the closure $\overline{R(L)}$ of $R(L)$.

Theorem 2: If the conditions C_1 - C_2 are satisfying ,then for any $\mathcal{F} = (f, \varphi_1, \varphi_2) \in H$, there exists a strong unique solution $u = \bar{L}^{-1}\mathcal{F} = \overline{\bar{L}^{-1}}\mathcal{F}$ of the probleme (1)-(5) where the estimate $\|u\|_B \leq c \|\mathcal{F}\|_H$ is satisfying ,where c is a positive constant does not depends of u .

Proof: From(22) we conclude that the operator \bar{L} acting from $D(\bar{L})$ in $R(\bar{L})$ have an inverse \bar{L}^{-1} , and from corollary 2, we conclude that the range $R(\bar{L})$ of the operator \bar{L} is closed. Then we will be prove the density of the set $R(L)$ in the space H (i.e) $\overline{R(L)} = H$.

For this we need the following proposition :

Proposition 2: If, for all functions $u \in D_0(L)$, where

$$D_0(L) = \{u/u \in D(L) : l_1 u = l_2 u = 0\}$$

and for some function $\omega \in L^2(Q)$, we have

$$(18) \quad (\mathcal{L}u, \omega)_{L^2(Q)} = 0,$$

then ω vanishes almost everywhere in Q .

Proof of the proposition 2 : The relation (18) is given for all $u \in D_0(L)$, we can express it in a particular form. Let u_{tt} be a solution of :

$$(19) \quad b(\sigma, t) [x \mathfrak{S}_x^* u_{tt} - \mathfrak{S}_x^*(\rho u_{tt})] = h(x, t),$$

where σ is a constant in $(0, l)$ and $h(x, t) = \int_t^T \omega(x, \tau) d\tau$. And let u be the function defined by:

$$(20) \quad u = \begin{cases} 0, & \text{si } 0 \leq t \leq s, \\ \int_s^t (t - \tau) u_{\tau\tau} d\tau, & \text{si } s \leq t \leq T. \end{cases}$$

(19) and (20) follows u is in $D_0(L)$ and:

$$(21) \quad \begin{aligned} \omega(x, t) &= \mathfrak{S}_x^{*-1} h \\ &= -[b(\sigma, t) (x \mathfrak{S}_x^* u_{tt} - \mathfrak{S}_x^*(\rho u_{tt}))]_t \\ &= [b(\sigma, t) \mathfrak{S}_x^*(\rho - x) u_{tt}]_t. \end{aligned}$$

To continue the proof we need the following lemma :

Lemma 2. The function ω defined by (21), belongs to the space $L^2(Q)$.

Proof of lemma 2 : We start with the proof of this inequality $\|\mathfrak{S}_x^*(\rho - x) u_{tt}\|_{L^2(0, l)}^2 \leq \frac{l^4}{12} \|u_{tt}\|_{L^2(0, 1)}^2$.

From this inequality and since the conditions C_1 are satisfied we conclude that $b_t(\sigma, t) \mathfrak{S}_x^*(\rho - x) u_{tt}$ belongs to $L^2(Q)$.

Because $\omega(x, t) = [b(\sigma, t) \mathfrak{S}_x^*(\rho - x) u_{tt}]_t = b_t(\sigma, t) \mathfrak{S}_x^*(\rho - x) u_{tt} + b(\sigma, t) \mathfrak{S}_x^*(\rho - x) u_{ttt}$, then we will

be proved that: $b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{ttt} \in L^2(Q)$.

For this we introduce the t -averaging operators ρ_ε of the form

$$(\rho_\varepsilon f)(x, t) = \frac{1}{\varepsilon} \int_0^T \omega\left(\frac{t-s}{\varepsilon}\right) f(x, s) ds,$$

where $\omega \in C_0^\infty(0, T)$, $\omega \geq 0$,

$\int_{-\infty}^{+\infty} \omega(s) ds = 1$, $\omega \equiv 0$ for $t \leq 0$ and $t \geq T$,

applying the operators ρ_ε and $\frac{\partial}{\partial t}$ to the equation

$$-b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt} = h(x, t),$$

we obtain

$$\begin{aligned} & \frac{\partial}{\partial t} (-b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt}) = \\ & \frac{\partial}{\partial t} [-b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt} + \rho_\varepsilon (b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt})] - \frac{\partial}{\partial t} \rho_\varepsilon h. \end{aligned}$$

Then

$$\begin{aligned} & \|b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt}\|_{L^2(Q)}^2 \\ & \leq 2 \left\| \frac{\partial}{\partial t} [b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt} - \rho_\varepsilon (b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt})] \right\|_{L^2(Q)}^2 \\ & \quad + 2 \left\| \frac{\partial}{\partial t} \rho_\varepsilon h \right\|_{L^2(Q)}^2. \end{aligned}$$

Since $\rho_\varepsilon f \xrightarrow{\varepsilon \rightarrow 0} f$, and $\frac{\partial}{\partial t} (b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt})$ is bounded in $L^2(Q)$, then $\omega \in L^2(Q)$.

Now we return to the 2nd proposition, we remplace ω in

(18) by its representation given by (21) we have:

$$\begin{aligned} (22) \quad & (u_{tt}, [b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt}]_t)_{L^2(Q)} \\ & = ((b(x, t)u_x)_x, [b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt}]_t)_{L^2(Q)} + \\ & \quad + \beta (u_{ttxx}, [b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt}]_t)_{L^2(Q)}. \end{aligned}$$

Making use conditions(3)-(5),and from the particular forme of u given by (19) and (20), the equality (22) can be simplified.For this integrating by parts each term of the equality on the sup-domain $Q_s = (0, l) \times (s, T)$ where $0 \leq s \leq T$

$$(23) \quad (u_{tt}, [b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt}]_t)_{L^2(Q)} \\ = \frac{1}{2} \left\| \sqrt{b(\sigma, s)} \mathfrak{S}_x^* u_{tt}(\cdot, s) \right\|_{L^2(0, l)}^2 - \frac{1}{2} \left\| \sqrt{b_t(\sigma, \cdot)} \mathfrak{S}_x^* u_{tt} \right\|_{L^2(Q_s)}^2,$$

$$(24) \quad ((b(x, t) u_x)_x, [b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt}]_t)_{L^2(Q)} \\ = -\frac{1}{2} \left\| \sqrt{b(\cdot, T) b(\sigma, T)} u_t(\cdot, T) \right\|_{L^2(0, l)}^2 \\ + \frac{1}{2} \int_{Q_s} [3b_t(x, t) b(\sigma, t) + b(x, t) b_t(\sigma, t)] (u_t)^2 dx dt \\ - \int_0^l b_t(x, T) b(\sigma, T) u(x, T) u_t(x, T) dx \\ + \int_{Q_s} [b_{tt}(x, t) b(\sigma, t) + b_t(x, t) b_t(\sigma, t)] u u_t dx dt \\ + \int_{Q_s} [b_x(x, t) u_t + b_{xt}(x, t) u] b(\sigma, t) \mathfrak{S}_x^* u_{tt} dx dt,$$

$$(25) \quad \beta (u_{ttxx}, [b(\sigma, t) \mathfrak{S}_x^*(\rho - x)u_{tt}]_t)_{L^2(Q)} \\ = \frac{\beta}{2} \left\| \sqrt{b_t(\sigma, \cdot)} u_{tt} \right\|_{L^2(Q_s)}^2 - \frac{\beta}{2} \left\| \sqrt{b(\sigma, s)} u_{tt}(\cdot, s) \right\|_{L^2(0, l)}^2.$$

Substitution of (23)-(25) into (22) gives

$$\begin{aligned}
(26) \quad & \frac{1}{2} \left\| \sqrt{b(\sigma, s)} \mathfrak{S}_x^* u_{tt}(\cdot, s) \right\|_{L^2(0, l)}^2 \\
& + \frac{1}{2} \left\| \sqrt{b(\cdot, T) b(\sigma, T)} u_t(\cdot, T) \right\|_{L^2(0, l)}^2 \\
& + \frac{\beta}{2} \left\| \sqrt{b(\sigma, s)} u_{tt}(\cdot, s) \right\|_{L^2(0, l)}^2 \\
= & \frac{1}{2} \left\| \sqrt{b_t(\sigma, s)} \mathfrak{S}_x^* u_{tt} \right\|_{L^2(Q_s)}^2 + \frac{\beta}{2} \left\| \sqrt{b_t(\sigma, \cdot)} u_{tt} \right\|_{L^2(Q_s)}^2 \\
& + \frac{1}{2} \int_{Q_s} [3b_t(x, t) b(\sigma, t) + b(x, t) b_t(\sigma, t)] (u_t)^2 dx dt \\
& - \int_0^l b_t(x, T) b(\sigma, T) u(x, T) u_t(x, T) dx \\
& + \int_{Q_s} [b_{tt}(x, t) b(\sigma, t) + b_t(x, t) b_t(\sigma, t)] u u_t dx dt \\
& + \int_{Q_s} [b_x(x, t) u_t + b_{xt}(x, t) u] b(\sigma, t) \mathfrak{S}_x^* u_{tt} dx dt .
\end{aligned}$$

By applying the Cauchy inequality and Cauchy inequality with ε to estimate the last three terms on the right-hand side of the inequality (26) and making use conditions $C_1 - C_2$, combining the estimates and (26) taking into account

that $\varepsilon = \frac{b_0^2}{2b_1^2}$ we obtain

$$\begin{aligned}
 (27) \quad & \frac{b_0}{2} \left[\|\mathfrak{S}_x^* u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 + \frac{b_0}{2} \|u_t(\cdot, T)\|_{L^2(0,l)}^2 + \right. \\
 & \quad \left. + \beta \|u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 \right] \\
 & \leq \left(b_1^2 + \frac{b_2}{2} \right) \|\mathfrak{S}_x^* u_{tt}\|_{L^2(Q_s)}^2 + \frac{\beta b_2}{2} \|u_{tt}\|_{L^2(Q_s)}^2 + \\
 & \quad + \frac{b_2^2 + b_1^2 + 4b_1 b_2 + b_3^2}{2} \|u_t\|_{L^2(Q_s)}^2 + \\
 & \quad + \frac{b_2^2 + b_4^2 + b_5^2}{2} \|u\|_{L^2(Q_s)}^2 + \frac{b_1^2 b_2^2}{b_0^2} \|u(\cdot, T)\|_{L^2(0,l)}^2
 \end{aligned}$$

By virtue of the elementary inequality

$$(28) \quad \frac{b_1^2 b_2^2}{b_0^2} \|u(\cdot, T)\|_{L^2(0,l)}^2 \leq \frac{b_1^2 b_2^2}{b_0^2} \|u\|_{L^2(Q_s)}^2 + \frac{b_1^2 b_2^2}{b_0^2} \|u_t\|_{L^2(Q_s)}^2,$$

we estimate the last term of the right-hand side of the inequality(27),we obtain

$$\begin{aligned}
 (29) \quad & \|\mathfrak{S}_x^* u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 + \\
 & + \frac{b_0}{2} \|u_t(\cdot, T)\|_{L^2(0,l)}^2 + \beta \|u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 \\
 & \leq \left(\frac{2b_1^2 + b_2}{b_0} \right) \|\mathfrak{S}_x^* u_{tt}\|_{L^2(Q_s)}^2 + \frac{\beta b_2}{b_0} \|u_{tt}\|_{L^2(Q_s)}^2 \\
 & \quad + \frac{\frac{2b_1^2 b_2^2}{b_0^2} + 2b_1 b_2 + (b_2 + b_1)^2 + b_3^2}{b_0} \|u_t\|_{L^2(Q_s)}^2 \\
 & \quad + \frac{b_0^2 (b_2^2 + b_4^2 + b_5^2) + 2b_1^2 b_2^2}{b_0^3} \|u\|_{L^2(Q_s)}^2
 \end{aligned}$$

For estimate the last term of the right-hand side of the inequality(29),we will be proove the inequality $\|u\|_{L^2(Q_s)}^2 \leq 24T^2 \|u_t\|_{L^2(Q_s)}^2$, combining the last inequality and (29) we get

$$\begin{aligned}
(30) \quad & \|\mathfrak{S}_x^* u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 + \|u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 + \\
& + \|u_t(\cdot, T)\|_{L^2(0,l)}^2 \\
& \leq k \left[\|\mathfrak{S}_x^* u_{tt}\|_{L^2(Q_s)}^2 + \|u_{tt}\|_{L^2(Q_s)}^2 + \|u_t\|_{L^2(Q_s)}^2 \right],
\end{aligned}$$

$$\text{where } k = \frac{\max(\beta b_2, (2b_1^2 + b_2), b_0 k(b_i, T))}{b_0 \min(1, \beta, \frac{b_0}{2})}.$$

$$k(b_i, T) = \frac{2b_1^2 b_2^2 + b_0^2 [2b_1 b_2 + (b_2 + b_1)^2 + b_3^2] + 24T^2 [b_0^2 (b_2^2 + b_4^2 + b_5^2) + 2b_1^2 b_2^2]}{b_0^3}.$$

To continue, we introduce the new function $v(x, t) = \int_t^T u_{\tau\tau} d\tau$, then $u_t(x, t) = v(x, s) - v(x, t)$, and $u_t(x, T) = v(x, s)$.

The inequality (30) it be

$$\begin{aligned}
(31) \quad & \|\mathfrak{S}_x^* u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 + \|u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 + \\
& + (1 - 2k(T - s)) \|v(\cdot, s)\|_{L^2(0,l)}^2 \\
& \leq 2k \left(\|\mathfrak{S}_x^* u_{tt}\|_{L^2(Q_s)}^2 + \|u_{tt}\|_{L^2(Q_s)}^2 + \|v\|_{L^2(Q_s)}^2 \right).
\end{aligned}$$

If $s_0 > 0$ satisfies $(1 - 2k(T - s_0)) = \frac{1}{2}$, then the inequality (31) implies

$$\begin{aligned}
(32) \quad & \|\mathfrak{S}_x^* u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 + \|u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 \\
& + \|v(\cdot, s)\|_{L^2(0,l)}^2 \\
& \leq 4k \left(\|\mathfrak{S}_x^* u_{tt}\|_{L^2(Q_s)}^2 + \|u_{tt}\|_{L^2(Q_s)}^2 + \|v\|_{L^2(Q_s)}^2 \right),
\end{aligned}$$

for all $s \in [T - s_0, T]$. We denote

$$Y(s) = \|\mathfrak{S}_x^* u_{tt}\|_{L^2(Q_s)}^2 + \|u_{tt}\|_{L^2(Q_s)}^2 + \|v\|_{L^2(Q_s)}^2.$$

We get:

$$Y'(s) = -\|\mathfrak{S}_x^* u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 - \|u_{tt}(\cdot, s)\|_{L^2(0,l)}^2 - \|v(\cdot, s)\|_{L^2(0,l)}^2.$$

Then and from (32) we obtain $-Y'(s) \leq 4kY(s)$.
 Then $-\frac{\partial}{\partial s}(Y(s) \exp(4ks)) \leq 0$.

Integrating this inequality on (s, T) and taking into account that $Y(T) = 0$, we obtain $Y(s) \exp(4ks) \leq 0$.
 Then $Y(s) = 0$ for all $s \in [T - s_0, T]$. Then $\omega = 0$ almost everywhere in Q_{T-s_0} , proceeding in this way step by step, we prove that $\omega = 0$ almost everywhere in Q .

This achieves the proof of proposition. Now we return to prove the théorème. We will prove that $\overline{R(L)} = H$.
 Since H is a Hilbert space, the equality $\overline{R(L)} = H$ is true, if from

$$(33) \quad (Lu, W)_H = (\mathcal{L}u, \omega)_{L^2(Q)} + (l_1u, \omega_1)_{L^2(0,l)} + (l_2u, \omega_2)_{L^2(0,l)} = 0,$$

where $W = (\omega, \omega_1, \omega_2) \in R(L)^\perp$, we get $\omega \equiv 0, \omega_1 \equiv 0$ and $\omega_2 \equiv 0$ in Q , for any element of $D_0(L)$.

From (33) we obtain $\forall u \in D_0(L), (\mathcal{L}u, \omega)_{L^2(Q)} = 0$. Then by virtue of the 2nd proposition, we conclude that $\omega \equiv 0$.

Then for (33), we obtain $(l_1u, \omega_1)_{L^2(0,l)} + (l_2u, \omega_2)_{L^2(0,l)} = 0$.

Since the quantities l_1u and l_2u can vanish independently and the ranges of the trace operators l_1 and l_2 are dense in the Hilbert space $L^2(0, l)$, then $\omega_1 = \omega_2 = 0$. Thus to conclude that $W = 0$.

REFERENCES

- [1] A. BOUZIANI, *Solution forte d'un problème mixte avec conditions non locales pour une classe d'équations hyperboliques*, Bulletin de la Classe des Sciences, Académie Royale de Belgique, 1-6, (1997).
- [2] A. BOUZIANI, *Mixed problem with boundary integral conditions for a certain parabolic equation*, J. Appl. Math and Stochastic Analysis, Vol. 9, N°3, 323-330, (1996).
- [3] A. BOUZIANI, *Problèmes mixtes avec conditions intégrales pour quelques équations aux dérivées partielles*, Thèse de doctorat d'Etat, Université de constantine, Algeria (1996).
- [4] A. BOUZIANI – N. E. BENOUAR, *Problème mixte avec conditions intégrales pour une classe d'équations paraboliques*, Comptes Rendus de l'Académie des Sciences. Paris, Vol. 321, Série I, 1177-1182, (1995).
- [5] J. R. CANNON, *The solution of the heat equation subject to the specification of energy*, Quart. Appl. Math., Vol. 21, N°2, 155-160, (1963).
- [6] A. A. DEZIN, *Théorèmes d'existence et d'unicité de la solution pour les problèmes aux limites des équations aux dérivées partielles dans les espaces fonctionnels*, Uspekhi. Math. Naouk, 14, N°3 (87), 22-73, (1959).
- [7] L. GARDING, *Cauchy's problem for hyperbolic equations*, University of Chicago, Lecture notes, (1957).
- [8] O. A. LADYZHENSKAYA, *Sur les problèmes aux limites fondamentaux liés aux équations paraboliques et hyperboliques*, Dokl. Acad. Scien. URSS, Vol. 97, N°3, 395-398, (1954).
- [9] P. LAX, *On Cauchy's problem for hyperbolic equation and the differentiability of solutions of elliptic equations*. Comm. Pure and Appl. Math., Vol. 8, 615-633, (1955).
- [10] J. LERAY, *Lectures on hyperbolic differential equations with variable coefficients*. Princeton, Just for Adv. study, (1952).
- [11] S. MESLOUB, *Problème mixtes avec conditions non locales pour certaines classes d'équations aux dérivées partielles*, Thèse de doctorat d'Etat, Université de constantine, (1999).
- [12] R. SAKOMOTO, *Mixed problems for hyperbolic equations II*, Jour. Math. Kyoto Univ., 10, 3, 403-417, (1970).

1.KING SAOUD UNIVERSITY,DEPARTEMENT OF MATHEMATICS RIADH,ARABIE SAOUDITE.

2.INSTITUT CAMILLE JORDAN,UNIVERSIT LYON1,FRANCE.

E-mail address: 1. mesloub@yahoo.com ,2. amansour@math.univ-lyon1.fr